

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 952 221 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

27.10.1999 Bulletin 1999/43

(51) Int Cl.⁶: C12N 15/53, C12N 15/60,

C12N 1/21, C12N 1/20,

C12P 13/14

// (C12N1/20, C12R1:01, 1:425)

(21) Application number: 99105508.8

(22) Date of filing: 17.03.1999

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 18.03.1998 JP 6906898

19.10.1998 JP 29712998

(71) Applicant: Ajinomoto Co., Ltd.
Tokyo (JP)

(72) Inventors:

- Izui, Hiroshi c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)

- Ono, Eiji c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Matsui, Kazuhiko c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Moriya, Mika c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Ito, Hisao c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Hara, Yoshihiko c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)

(74) Representative: Strehl Schübel-Hopf & Partner
Maximilianstrasse 54
80538 München (DE)

(54) **L-glutamic acid-producing bacterium and method for producing L-glutamic acid**

(57) L-Glutamic acid is produced by culturing in a liquid culture medium a microorganism belonging to the genus *Enterobacter* or *Serratia* and having an ability to produce L-glutamic acid, which increases in an activity of enzyme catalyzing a reaction for L-glutamic acid bio-

synthesis, or which decreases in or is deficient in an activity of an enzyme catalyzing a reaction branching from a pathway for L-glutamic acid biosynthesis and producing a compound other than L-glutamic acid, and collecting produced L-glutamic acid from the culture medium.

Description

BACKGROUND OF THE INVENTION

[0001] The present invention relates to a novel L-glutamic acid-producing bacterium and a method for producing L-glutamic acid by fermentation using the same. L-Glutamic acid is an important amino acid as food, drugs and the like.

[0002] L-Glutamic acid has conventionally been produced by fermentation methods utilizing the so-called coryneform L-glutamic acid-producing bacteria which principally belong to the genera *Brevibacterium*, *Corynebacterium*, and *Microbacterium* or variants thereof ("Amino Acid Fermentation", Gakkai Shuppan Center, pp.195-215, 1986). As methods for producing L-glutamic acid by fermentation utilizing other bacterial strains, there have been known the methods utilizing microorganisms of the genera *Bacillus*, *Streptomyces*, *Penicillium* and the like (United States Patent No. 3,220,929), the methods utilizing microorganisms of the genera *Pseudomonas*, *Arthrobacter*, *Serratia*, *Candida* and the like (United States Patent No. 3,563,857), the methods utilizing microorganisms of the genera *Bacillus*, *Pseudomonas*, *Serratia* and the like or *Aerobacter aerogenes* (currently referred to as *Enterobacter aerogenes*) (Japanese Patent Publication (KOKOKU) No. 32-9393(1957)), the method utilizing variant strains of *Escherichia coli* (Japanese Patent Application Laid-Open (KOKAI) No. 5-244970(1993)) and the like.

[0003] Though the productivity of L-glutamic acid has considerably been improved by breeding of such microorganisms as mentioned above or improvements of production methods, it is still desired to develop a more inexpensive and more efficient method for producing L-glutamic acid in order to meet the expected markedly increasing future demand of the amino acid.

SUMMARY OF THE INVENTION

[0004] The object of the present invention is to find a novel L-glutamic acid-producing bacterium having a high ability to produce L-glutamic acid, thereby developing a more inexpensive and more efficient method for producing L-glutamic acid.

[0005] To achieve the aforementioned object, the present inventors intensively searched for and studied microorganisms having the ability to produce L-glutamic acid that are different from the previously reported microorganisms. As a result, they found that certain strains derived from microorganisms belonging to the genus *Enterobacter* or *Serratia* had a high ability to produce L-glutamic acid, and have completed the present invention.

[0006] Thus, the present invention provides:

(1) a microorganism belonging to the genus *Enterobacter* or *Serratia* and having an ability to produce L-glutamic acid and at least one of the following properties:

- (a) the microorganism increases in an activity of an enzyme catalyzing a reaction for L-glutamic acid biosynthesis; and
- (b) the microorganism decreases in or is deficient in an activity of an enzyme catalyzing a reaction branching from a pathway for L-glutamic acid biosynthesis and producing a compound other than L-glutamic acid;

(2) a microorganism of the above (1) wherein the enzyme catalyzing the reaction for the L-glutamic acid biosynthesis is at least one selected from the group consisting of citrate synthase (abbreviated as "CS" hereinafter), phosphoenolpyruvate carboxylase (abbreviated as "PEPC" hereinafter), and glutamate dehydrogenase (abbreviated as "GDH" hereinafter);

(3) a microorganism of the above (2) wherein the enzyme catalyzing the reaction for the L-glutamic acid biosynthesis includes all of CS, PEPC and GDH;

(4) a microorganism of any one of the above (1) to (3) wherein the enzyme catalyzing the reaction branching from the pathway for L-glutamic acid biosynthesis and producing the compound other than L-glutamic acid is α -ketoglutarate dehydrogenase (abbreviated as "aKGDH" hereinafter);

(5) a microorganism of any one of the above (1) to (4) which is *Enterobacter agglomerans* or *Serratia liquefacience*; and

(6) a method for producing L-glutamic acid which comprises culturing the microorganism as defined in any one of the above (1) to (5) in a liquid culture medium to produce and accumulate L-glutamic acid in the culture medium, and collecting the L-glutamic acid from the culture medium.

[0007] Because the microorganism of the present invention have a high ability to produce L-glutamic acid, it is considered to be possible to impart a further higher production ability to the microorganism by using the breeding techniques previously known for the coryneform L-glutamic acid-producing bacteria and the like, and it is expected to contribute

to development of a more inexpensive and more efficient method for producing L-glutamic acid by appropriately selecting culture conditions and the like.

BRIEF EXPLANATION OF THE DRAWINGS

[0008] Figure 1 shows construction of a plasmid pMWCPG having a *gltA* gene, a *ppc* gene and a *gdhA* gene.
 [0009] Figure 2 shows construction of a plasmid pSTVG having the *gdhA* gene.
 [0010] Figure 3 shows construction of a plasmid RSF-Tet having a replication origin of a wide-host-range plasmid RSF1010 and a tetracycline resistance gene.
 [0011] Figure 4 shows construction of a plasmid RSFCPG having the replication origin of the wide-host-range plasmid RSF1010, the tetracycline resistance gene, the *gltA* gene, the *ppc* gene and the *gdhA* gene.
 [0012] Figure 5 shows construction of a plasmid pMWCB having the *gltA* gene.
 [0013] Figure 6 shows a restriction map of a DNA fragment of pTWVEK101 derived from *Enterobacter agglomerans*.
 [0014] Figure 7 shows comparison of an amino acid sequence deduced from a nucleotide sequence of a *sucA* gene derived from *Enterobacter agglomerans* with one derived from *Escherichia coli*. The upper sections: *Enterobacter agglomerans*, the lower sections: *Escherichia coli* (the same shall apply hereinafter).
 [0015] Figure 8 shows comparison of an amino acid sequence deduced from a nucleotide sequence of a *sucB* gene derived from *Enterobacter agglomerans* with one derived from *Escherichia coli*.
 [0016] Figure 9 shows comparison of an amino acid sequence deduced from a nucleotide sequence of a *sdhB* gene derived from *Enterobacter agglomerans* with one derived from *Escherichia coli*.
 [0017] Figure 10 shows comparison of an amino acid sequence deduced from a nucleotide sequence of a *sucC* gene derived from *Enterobacter agglomerans* with one derived from *Escherichia coli*.

DETAILED DESCRIPTION OF THE INVENTION

[0018] The present invention will be explained in detail hereinafter.
 [0019] The microorganism of the present invention is a microorganism belonging to the genus *Enterobacter* or *Serratia*, and having at least one of the following properties:

- (a) the microorganism increases in an activity of an enzyme catalyzing a reaction for L-glutamic acid biosynthesis; and
- (b) the microorganism decreases in or is deficient in an activity of an enzyme catalyzing a reaction branching from a pathway for L-glutamic acid biosynthesis and producing a compound other than L-glutamic acid.

[0020] Such a microorganism can be obtained by using a microorganism belonging to the genus *Enterobacter* or the genus *Serratia* as a parent strain, and imparting the properties of the above (a) and/or (b) to the microorganism. Examples of the microorganism belonging to the genus *Enterobacter* or the genus *Serratia* that can be used as the parent strain are listed below:

Enterobacter agglomerans
Enterobacter aerogenes
Enterobacter amnigenus
Enterobacter asburiae
Enterobacter cloacae
Enterobacter dissolvens
Enterobacter gergoviae
Enterobacter hormaechei
Enterobacter intermedius
Enterobacter nimipressuralis
Enterobacter sakazakii
Enterobacter taylorae
Serratia liquefacience
Serratia entomophila
Serratia ficaria
Serratia fonticola
Serratia grimesii
Serratia proteamaculans
Serratia odorifera

Serratia plymuthica
Serratia rubidaea

[0021] More preferably, those bacterial strains listed below can be mentioned:

Enterobacter agglomerans ATCC 12287
Enterobacter agglomerans AJ13355
Serratia liquefacience ATCC 14460

[0022] The *Enterobacter agglomerans* AJ13355 was deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry on February 19, 1998, and received an accession number of FERM P-16644, and then transferred to an international deposition under the Budapest Treaty on January 11, 1999, and received an accession number of FERM BP-6614. The *Enterobacter agglomerans* ATCC 12287, and the *Serratia liquefacience* ATCC 14460 are available from ATCC.

[0023] The *Enterobacter agglomerans* AJ13355 is a strain isolated from soil in Iwata-shi, Shizuoka, Japan.

[0024] Physiological properties of AJ13355 are as follows:

- (1) Gram stain: Negative
- (2) Behavior for oxygen: Facultative anaerobe
- (3) Catalase: Negative
- (4) Oxidase: Positive
- (5) Nitrate reduction: Negative
- (6) Voges-Proskauer reaction: Positive
- (7) Methyl Red test: Negative
- (8) Urease: Negative
- (9) Indole production: Positive
- (10) Motility: Present
- (11) Hydrogen sulfide production in TSI culture medium: Slightly active
- (12) β -Galactosidase: Positive
- (13) Sugar assimilability:

Arabinose: Positive
Sucrose: Positive
Lactose: Positive
Xylose: Positive
Sorbitol: Positive
Inositol: Positive
Trehalose: Positive
Maltose: Positive
Melibiose: Positive
Adonitol: Negative
Raffinose: Positive
Salicin: Negative
Melibiose: Positive

- (14) Glycerose assimilability: Positive
- (15) Organic acid assimilability:

Citric acid: Positive
Tartaric acid: Negative
Gluconic acid: Positive
Acetic acid: Positive
Malonic acid: Negative

- (16) Arginine dehydratase: Negative
- (17) Ornithine decarboxylase: Negative
- (18) Lysine decarboxylase: Negative
- (19) Phenylalanine deaminase: Negative

(20) Pigment formation: Yellow

(21) Gelatin liquefaction: Positive

(22) Growth pH: Not good growth at pH 4, good growth at pH 4.5 to 7

(23) Growth temperature: Good growth at 25°C, good growth at 30°C, good growth at 37°C, growth possible at 42°C, no growth at 45°C

[0025] From these bacteriological properties, AJ13355 is determined to be *Enterobacter agglomerans*.

[0026] In the working examples described hereinafter, *Enterobacter agglomerans* ATCC12287, *Enterobacter agglomerans* AJ13355, and *Serratia liquefacience* ATCC14460 were used as starting parent strains for obtaining strains which increase in the activity of the enzyme catalyzing the reactions for the L-glutamic acid biosynthesis, or strains which decrease in or are deficient in the activity of the enzyme catalyzing the reaction branching from the pathway for L-glutamic acid biosynthesis and producing the compound other than L-glutamic acid. However, the sugar metabolism by any of bacteria belonging to the genera *Enterobacter* and *Serratia* is achieved via the Embden-Meyerhof pathway, and pyruvate produced in the pathway is oxidized in the tricarboxylic acid cycle under aerobic conditions. L-Glutamic acid is biosynthesized from α -ketoglutaric acid which is an intermediate of the tricarboxylic acid cycle by GDH or glutamine synthetase/glutamate synthase. Thus, these microorganisms share the same biosynthetic pathway for L-glutamic acid, and microorganism belonging to the genera *Enterobacter* and *Serratia* are encompassed within a single concept according to the present invention. Therefore, microorganisms belonging to the genera *Enterobacter* and *Serratia* other than species and strains specifically mentioned above also fall within the scope of the present invention.

[0027] The microorganism of the present invention is a microorganism belonging to the genus *Enterobacter* or the genus *Serratia* and having an ability to produce L-glutamic acid. The expression "having an ability to produce L-glutamic acid" as herein used means to have an ability to accumulate L-glutamic acid in a culture medium during cultivation. According to the present invention, the ability to produce L-glutamic acid is imparted by giving either one or both of the following characteristics:

- (a) the microorganism increases in the activity of the enzyme catalyzing the reaction for the L-glutamic acid biosynthesis; and
- (b) the microorganism decreases in or is deficient in the activity of the enzyme catalyzing the reaction branching from the pathway for L-glutamic acid biosynthesis and producing the compound other than L-glutamic acid.

[0028] As examples of the enzyme catalyzing the reaction for L-glutamic acid biosynthesis of microorganisms of the genus *Enterobacter* or *Serratia*, there can be mentioned GDH, glutamine synthetase, glutamate synthase, isocitrate dehydrogenase, aconitate hydratase, CS, PEPC, pyruvate dehydrogenase, pyruvate kinase, enolase, phosphoglyceromutase, phosphoglycerate kinase, glyceraldehyde-3-phosphate dehydrogenase, triosephosphate isomerase, fructose bisphosphate aldolase, phosphofructokinase, glucose phosphate isomerase and the like. Among these enzymes, one or two or three kinds of CS, PEPC and GDH are preferred. As for the microorganism of the present invention, it is further preferred that activities of all of the three kinds of enzymes, CS, PEPC and GDH, are increased. Whether a microorganism increases in an activity of a target enzyme, and degree of the increase of the activity can be determined by measuring the enzyme activity of a bacterial cell extract or a purified fraction, and comparing it with that of a wild type strain or a parent strain.

[0029] The microorganism of the present invention, which belongs to the genus *Enterobacter* or *Serratia*, and increases in the activity of the enzyme catalyzing the reaction for L-glutamic acid biosynthesis, can be obtained as, for example, a variant where mutation has been made in a gene encoding the enzyme or a genetic recombinant strain by using any of the microorganisms mentioned above as a starting parent strain.

[0030] To enhance the activity of CS, PEPC or GDH, for example, a gene encoding CS, PEPC or GDH can be cloned in a suitable plasmid, and the aforementioned starting parent strain as a host can be transformed with the resulting plasmid. This can increase the copy number of each of the genes encoding CS, PEPC and GDH (hereinafter abbreviated as "gltA gene", "ppc gene", and "gdhA gene", respectively), and as a result the activities of CS, PEPC and GDH can be increased.

[0031] One or two or three kinds selected from the cloned gltA gene, ppc gene and gdhA gene in any combination are introduced into the starting parent strain mentioned above. When two or three kinds of the genes are introduced, either the two or three kinds of the genes are cloned in one kind of plasmid, and introduced into the host, or they are separately cloned in two or three kinds of plasmids that can exist in the same host, and introduced into the host.

[0032] The plasmid is not particularly limited so long as it can autonomously replicate in a microorganism belonging to the genus *Enterobacter* or *Serratia*. Examples of the plasmid include, for example, pUC19, pUC18, pBR322, pHSG299, pHSG298, pHSG399, pHSG398, RSF1010, pMW119, pMW118, pMW219, pMW218 and the like. Other than these plasmids, phage DNA vectors can also be utilized.

[0033] Transformation can be achieved by, for example, the method of D.M. Morrison (Methods in Enzymology 68,

326 (1979)), the method by increasing permeability of recipient cells for DNA with calcium chloride (Mandel, M. and Higa, A., J. Mol. Biol., 53, 159 (1970)), or the like.

[0034] The activities of CS, PEPC and GDH can also be increased by using multiple copies of the *gltA* gene, the *ppc* gene and/or the *gdh* gene present on the chromosome DNA of the starting parent strain as a host. In order to introduce multiple copies of the *gltA* gene, the *ppc* gene and/or the *gdhA* gene into a chromosome DNA of a microorganism belonging to the genus *Enterobacter* or *Serratia*, sequences present on chromosome DNA in a multiple copy number such as repetitive DNA, and inverted repeats present at an end of transposition factors can be utilized. Alternatively, multiple copies of the genes can also be introduced into a chromosome DNA by utilizing transposition of transposons carrying the *gltA* gene, the *ppc* gene, or the *gdhA* gene. These techniques can increase the copy number of the *gltA* gene, the *ppc* gene, and the *gdhA* gene in transformant cells, and as a result increase the activities of CS, PEPC and GDH.

[0035] Any organisms can be used as a source of the *gltA* gene, the *ppc* gene and the *gdhA* gene used for increasing copy numbers, so long as the organisms have the CS, PEPC and GDH activities. Among such organisms, bacteria, i. e., prokaryotes, such as those bacteria belonging to the genera *Enterobacter*, *Klebsiella*, *Erwinia*, *Pantoea*, *Serratia*, *Escherichia*, *Corynebacterium*, *Brevibacterium*, and *Bacillus* are preferred. As a specific example, *Escherichia coli* can be mentioned. The *gltA* gene, the *ppc* gene and the *gdhA* gene can be obtained from a chromosome DNA of such microorganisms as mentioned above.

[0036] The *gltA* gene, the *ppc* gene and the *gdhA* gene can each be obtained from a chromosome DNA of any of the aforementioned microorganisms by isolating a DNA fragment complementing auxotrophy of a variant strain lacking the CS, PEPC or GDH activity. Alternatively, because the nucleotide sequences of these genes of bacteria of the genus *Escherichia* or *Corynebacterium* have already been elucidated (Biochemistry, Vol. 22, pp.5243-5249, 1983; J. Biochem. Vol. 95, pp.909-916, 1984; Gene, Vol. 27, pp.193-199, 1984; Microbiology, Vol. 140, pp.1817-1828, 1994; Mol. Gen. Genet. Vol. 218, pp.330-339, 1989; and Molecular Microbiology, Vol. 6, pp.317-326, 1992), the genes can be obtained by PCR using primers synthesized based on each of the elucidated nucleotide sequences, and the chromosome DNA as a template.

[0037] The activity of CS, PEPC or GDH can also be increased by, other than by the gene amplification mentioned above, enhancing expression of the *gltA* gene, the *ppc* gene or the *gdhA* gene. For example, the expression is enhanced by replacing the promoter of the *gltA* gene, the *ppc* gene, or the *gdhA* gene with another stronger promoter. Examples of such a strong promoter include, for example, a *lac* promoter, a *trp* promoter, a *trc* promoter, a *tac* promoter, a P_R promoter and a P_L promoter of lambda phage and the like. The *gltA* gene, the *ppc* gene, or the *gdhA* gene of which promoter has been substituted is cloned into a plasmid and introduced into a host microorganism, or introduced into a chromosome DNA of host microorganism using a repetitive DNA, inverted repeat, transposon or the like.

[0038] The activities of CS, PEPC or GDH can also be increased by replacing the promoter of the *gltA* gene, the *ppc* gene, or the *gdhA* gene on a chromosome with another stronger promoter (see WO87/03006, and Japanese Patent Application Laid-Open (KOKAI) No. 61-268183(1986)), or inserting a strong promoter at the upstream of each coding sequence of the genes (see Gene, 29, pp. 231-241, 1984). Specifically, these are achieved by homologous recombination between the *gltA* gene, the *ppc* gene, or the *gdhA* gene of which promoter is replaced with a stronger promoter or DNA containing a part of them, and a corresponding gene on the chromosome.

[0039] Specific examples of the microorganism belonging to the genus *Enterobacter* or *Serratia* of which CS, PEPC or GDH activity is increased include, for example, *Enterobacter agglomerans* ATCC12287/RSFCPG, *Enterobacter agglomerans* AJ13355/RSFCPG, and *Serratia liquefacience* ATCC14460/RSFCPG.

[0040] Examples of the enzyme catalyzing the reaction branching from the pathway of L-glutamic acid biosynthesis and producing the compound other than L-glutamic acid include, for example, α KGDH, isocitrate lyase, phosphate acetyltransferase, acetate kinase, acetohydroxy acid synthase, acetolactate synthase, formate acetyltransferase, lactate dehydrogenase, glutamate decarboxylase, 1-pyrroline dehydrogenase and the like. Among these enzymes, α KGDH is preferred.

[0041] In order to obtain such decrease or deficiency of enzyme activity as mentioned above in a microorganism belonging to the genus *Enterobacter* or *Serratia*, a mutation causing the decrease or deficiency of the enzyme activity can be introduced into a gene encoding the enzyme by a conventional mutagenesis technique or genetic engineering technique.

[0042] Examples of the mutagenesis technique include, for example, the method utilizing irradiation of X-ray or ultraviolet light, the method utilizing treatment with a mutagenic agent such as N-methyl-N'-nitro-N-nitrosoguanidine and the like. The site of gene to which a mutation is introduced may be a coding region encoding an enzyme protein, or an expression regulatory region such as a promoter.

[0043] Examples of the genetic engineering technique include, for example, genetic recombination, genetic transduction, cell fusion and the like. For example, a drug resistance gene is inserted into a target gene to produce a functionally inactivated gene (defective gene). Then, this defective gene is introduced into a cell of a microorganism belonging to the genus *Enterobacter* or *Serratia*, and the target gene on a chromosome is replaced with the defective

gene by homologous recombination (gene disruption).

[0044] Whether a microorganism decreases in an activity of a target enzyme or is deficient in the activity, and degree of the decrease of the activity can be determined by measuring the enzyme activity of a bacterial cell extract or a purified fraction of a candidate strain, and comparing it with that of a wild type strain or a parent strain. The α KGDH enzymatic activity can be measured by, for example, the method of Reed et al. (L.J. Reed and B.B. Mukherjee, Methods in Enzymology 1969, 13, p.55-61).

[0045] Depending on the target enzyme, a target variant can be selected based on a phenotype of the variant. For example, a variant which is deficient in the α KGDH activity or decreases in the activity cannot grow on a minimal medium containing glucose, or a minimal medium containing acetic acid or L-glutamic acid as an exclusive carbon source, or shows markedly reduced growth rate therein under aerobic conditions. However, even under the same condition, it can exhibit normal growth by addition of succinic acid or lysine, methionine and diaminopimelate to the minimal medium containing glucose. Based on these phenomena, a variant that is deficient in the α KGDH activity or decreases in the activity can be selected.

[0046] A method for producing a *Brevibacterium lactofermentum* strain lacking the α KGDH gene based on homologous recombination is detailed in WO95/34672, and a similar method can be used for microorganisms belonging to the genera *Enterobacter* and *Serratia*.

[0047] In addition, procedures of genetic cloning, cleavage and ligation of DNA, transformation and the like are detailed in Molecular Cloning, 2nd edition, Cold Spring Harbor Press (1989) and the like.

[0048] An example of the variant strain that is deficient in the α KGDH activity or decreases in the activity obtained as described above is *Enterobacter agglomerans* AJ13356. The *Enterobacter agglomerans* AJ13356 was deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry on February 19, 1998, received an accession number of FERM P-16645, and then transferred to an international deposition under the Budapest Treaty on January 11, 1999, and received an accession number of FERM BP-6615.

[0049] The microorganism belonging to the genus *Enterobacter* or *Serratia*, and having at least one of the properties (a) and (b) and an ability to produce L-glutamic acid can be cultured in a liquid medium to produce and accumulate L-glutamic acid in the medium.

[0050] The culture medium may be an ordinary nutrient medium containing a carbon source, a nitrogen source, and inorganic salts, as well as organic trace nutrients such as amino acids, vitamins and the like, as required. It can be a synthetic medium or a natural medium. Any carbon sources and nitrogen sources can be used for the culture medium so long as they can be utilized by the microorganism to be cultured.

[0051] The carbon source may be a saccharide such as glucose, glycerol, fructose, sucrose, maltose, mannose, galactose, starch hydrolysates, molasses and the like. Further, an organic acid such as acetic acid and citric acid may also be used alone or in combination with other carbon sources.

[0052] The nitrogen source may be ammonia, ammonium salts such as ammonium sulfate, ammonium carbonate, ammonium chloride, ammonium phosphate, and ammonium acetate, nitrates and the like.

[0053] As organic trace nutrients, amino acids, vitamins, fatty acids, nucleic acids, materials containing them such as peptone, casamino acid, yeast extract, and soybean protein decomposition products and the like are used, and when an auxotrophic variant which requires an amino acid or the like for its growth is used, it is necessary to complement the nutrient required.

[0054] As the inorganic salt, phosphates, magnesium salts, calcium salts, iron salts, manganese salts and the like are used.

[0055] As for the culture conditions, cultivation may be performed under aerobic conditions at a temperature of 20 to 42°C and a pH of 4 to 8. The cultivation can be continued for 10 hours to 4 days to accumulate a considerable amount of L-glutamic acid in the liquid culture medium.

[0056] After the completion of the cultivation, L-glutamic acid accumulated in the culture medium may be collected by a known method. For example, it can be isolated by a method comprising concentrating the medium after removing the cells to crystallize the product, ion exchange chromatography or the like.

Examples

[0057] The present invention will be explained more specifically with reference to the following examples.

(1) Construction of plasmid having *gltA* gene, *ppc* gene and *gdhA* gene

[0058] Procedure for construction of a plasmid having a *gltA* gene, a *ppc* gene and a *gdhA* gene will be explained by referring to Figure 1 to Figure 5.

[0059] A plasmid pBRGDH having a *gdhA* gene derived from *Escherichia coli* (Japanese Patent Application Laid-

Open (KOKAI) No. 7-203980(1995)) was digested with *HindIII* and *SphI*, and the both ends were blunt-ended by a treatment with T4 DNA polymerase. Then, a DNA fragment containing the *gdhA* gene was purified and collected. On the other hand, a plasmid pMWCP having a *gltA* gene and a *ppc* gene derived from *Escherichia coli* (WO97/08294) was digested with *XbaI*, and the both ends were blunt-ended by a treatment with T4 DNA polymerase. This was mixed with the DNA fragment having the *gdhA* gene purified above, and ligated with T4 ligase, giving a plasmid pMWCPG, which corresponds to the pMWCP further carrying the *gdhA* gene (Figure 1).

[0060] A DNA fragment having the *gdhA* gene obtained by digesting the pBRGDH with *HindIII* and *Sall* was purified and collected, and introduced into the *HindIII-Sall* site of a plasmid pSTV29 (purchased from Takara Shuzo) to obtain a plasmid pCTVG (Figure 2).

[0061] At the same time, a product obtained by digesting a plasmid pVIC40 having a replication origin of a wide-host-range plasmid RSF1010 (Japanese Patent Application Laid-Open (KOKAI) No. 8-047397(1996)) with *NotI*, followed by T4 DNA polymerase treatment and *PstI* digestion, and a product obtained by digesting pBR322 with *EcoT141*, followed by T4 DNA polymerase treatment and *PstI* digestion, were mixed and ligated with T4 ligase to obtain a plasmid RSF-Tet having the replication origin of RSF1010 and a tetracycline resistance gene (Figure 3).

[0062] Then, the pMWCPG was digested with *EcoRI* and *PstI*, and a DNA fragment having the *gltA* gene, the *ppc* gene and the *gdhA* gene was purified and collected. Similarly, the RSF-Tet was digested with *EcoRI* and *PstI*, and a DNA fragment having the replication origin of RSF1010 was purified and collected. Those DNA fragments were mixed and ligated with T4 ligase to obtain a plasmid RSFCPG composed of RSF-Tet carrying the *gltA* gene, the *ppc* gene and the *gdhA* gene (Figure 4). Expression of the *gltA* gene, the *ppc* gene and the *gdhA* gene by the resulting plasmid RSFCPG, and expression of the *gdhA* gene by the pSTVG were confirmed based on complementation of auxotrophy of *Escherichia coli* strains lacking the *gltA* gene, the *ppc* gene or the *gdhA* gene, and measurement of each enzyme activity.

[0063] A plasmid having a *gltA* gene derived from *Brevibacterium lactofermentum* was constructed as follows. PCR was performed by using primers having the nucleotide sequences represented in SEQ ID NOS: 6 and 7 selected based on the nucleotide sequence of the *gltA* gene of *Corynebacterium glutamicum* (Microbiology, 140, 1817-1828, 1994), and a chromosome DNA of *Brevibacterium lactofermentum* ATCC 13869 as a template to obtain a *gltA* gene fragment of about 3 kb. This fragment was inserted into a plasmid pHSG399 (purchased from Takara Shuzo) digested with *SmaI* to obtain a plasmid pHSGCB (Figure 5). Then, the pHSGCB was digested with *HindIII*, and an excised *gltA* gene fragment of about 3 kb was inserted into a plasmid pMW218 (purchased from Nippon Gene) digested with *HindIII* to obtain a plasmid pMWCB (Figure 5). Expression of the *gltA* gene by the resulting plasmid pMWCB was confirmed by determination of enzyme activity in the *Enterobacter agglomerans* AJ13355.

(2) Introduction of RSFCPG, pMWCB and pSTVG into *Enterobacter agglomerans* or *Serratia liquefacience*, and evaluation of L-glutamic acid productivity

[0064] The *Enterobacter agglomerans* ATCC 12287, the *Enterobacter agglomerans* AJ13355 and the *Serratia liquefacience* ATCC 14460 were transformed with the RSFCPG, pMWCB and pSTVG by electroporation (Miller J.H., "A Short Course in Bacterial Genetics; Handbook" Cold Spring Harbor Laboratory Press, USA, 1992) to obtain transformants exhibiting tetracycline resistance.

[0065] Each of the resulting transformants and the parent strains was inoculated into 50 ml-volume large size test tube containing 5 ml of a culture medium comprising 40 g/L glucose, 20 g/L ammonium sulfate, 0.5 g/L magnesium sulfate heptahydrate, 2 g/L potassium dihydrogenphosphate, 0.5 g/L sodium chloride, 0.25 g/L calcium chloride heptahydrate, 0.02 g/L ferrous sulfate heptahydrate, 0.02 g/L manganese sulfate tetrahydrate, 0.72 mg/L zinc sulfate dihydrate, 0.64 mg/L copper sulfate pentahydrate, 0.72 mg/L cobalt chloride hexahydrate, 0.4 mg/L boric acid, 1.2 mg/L sodium molybdate dihydrate, 2 g/L yeast extract, and 30 g/L calcium carbonate, and cultured at 37°C with shaking until the glucose contained in the culture medium was consumed. However, as for the AJ13355/pMWCB strain and the AJ13355/pSTVG strain, the cultivation was stopped when about 10 g/L of glucose was consumed, i.e., cultivated for 15 hours like the parent strain AJ13355, because their glucose consumption rates were low. To the culture medium of the transformants, 25 mg/L of tetracycline was added. After the cultivation was completed, L-glutamic acid accumulated in the culture medium was measured. The results are shown in Table 1.

Table 1:

Accumulated amount of L-glutamic acid	
Bacterial strain	Accumulated amount of L-glutamic acid
ATCC12287	0.0 g/L
ATCC12287/RSFCPG	6.1

Table 1: (continued)

Accumulated amount of L-glutamic acid	
Bacterial strain	Accumulated amount of L-glutamic acid
AJ13355	0.0
AJ13355/RSFCPG	3.3
AJ13355/pMWCB	0.8
AJ13355/pSTVG	0.8
ATCC14460	0.0
ATCC14460/RSFCPG	2.8
Culture medium alone	0.2

[0066] While the *Enterobacter agglomerans* ATCC12287, the *Enterobacter agglomerans* AJ13355 and the *Serratia liquefaciens* ATCC14460 did not accumulate L-glutamic acid, the strains whose CS, PEPC and GDH activities were amplified by introducing RSFCPG accumulated 6.1 g/L, 3.3 g/L, and 2.8 g/L of L-glutamic acid, respectively. The AJ13355 strain of which CS activity alone was amplified accumulated 0.8 g/L of L-glutamic acid, and the strain of which GDH activity alone was amplified also accumulated 0.8 g/L of L-glutamic acid.

(3) Cloning of α KGDH gene (referred to as "sucAB" hereinafter) of *Enterobacter agglomerans* AJ13355

[0067] The sucAB gene of the *Enterobacter agglomerans* AJ13355 was cloned by selecting a DNA fragment complementing acetate non-assimilation of an *Escherichia coli* strain lacking the α KGDH-E1 subunit gene (referred to as "sucA" hereinafter) from the chromosome DNA of the *Enterobacter agglomerans* AJ13355.

[0068] The chromosome DNA of the *Enterobacter agglomerans* AJ13355 strain was isolated by the same method as conventionally used for extracting chromosome DNA from *Escherichia coli* (Seibutsu Kogaku Jikken-sho (Textbook of Bioengineering Experiments), Ed. by the Society of Fermentation and Bioengineering, Japan, p.97-98, Baifukan, 1992). The pTWV228 used as the vector (ampicillin resistant) was a marketed product from Takara Shuzo.

[0069] Products obtained by digesting the chromosome DNA of the AJ13355 strain with *EcoT221* and products obtained by digesting the pTWV228 with *PstI* were ligated by T4 ligase, and the *Escherichia coli* JRG465 lacking sucA (Herbert J. et al., Mol. Gen. Genetics, 1969, 105, p.182) was transformed with them. Strains grown on the acetic acid minimal medium were selected from the transformants obtained as described above, and a plasmid extracted from them was designated as pTWVEK101. The *Escherichia coli* JRG465 carrying the pTWVEK101 recovered the characteristics of acetate non-assimilability as well as auxotrophy for succinic acid or L-lysine and L-methionine. This suggests that the pTWVEK101 contains the sucA gene of *Enterobacter agglomerans*.

[0070] A restriction map of *Enterobacter agglomerans*-derived DNA fragment of pTWVEK101 is shown in Figure 6. The result of nucleotide sequencing of the hatched portion in Figure 6 is shown in SEQ ID NO: 1. In this sequence, two full length ORFs and two nucleotide sequences considered as partial sequences of ORFs were found. Amino acid sequences that can be encoded by these ORFs and the partial sequences thereof are shown in SEQ ID NOS: 2 to 5 in order from the 5' ends. As a result of homology analysis of these sequences, it was found that the portion of which nucleotide sequence had been determined contained a 3' partial sequence of succinate dehydrogenase iron-sulfur protein gene (*sdhB*), full length sucA and α KGDH-E2 subunit gene (*sucB* gene), and 5' partial sequence of succinyl-CoA synthetase β subunit gene (*sucC* gene). Comparison of the amino acid sequences deduced from these nucleotide sequences with those of *Escherichia coli*. (Eur. J. Biochem., 141, 351-359 (1984), Eur. J. Biochem., 141, 361-374 (1984), and Biochemistry, 24, 6245-6252 (1985)) is shown in Figures 7 to 9. As shown by these results, the amino acid sequences exhibited markedly high homology. It was also found that a cluster of *sdhB-sucA-sucB-sucC* is formed on the *Enterobacter agglomerans* chromosome like *Escherichia coli* (Eur. J. Biochem., 141, 351-359 (1984), Eur. J. Biochem., 141, 361-374 (1984), and Biochemistry, 24, 6245-6252 (1985)).

(4) Acquisition of strain deficient in α KGDH derived from *Enterobacter agglomerans* AJ13355

[0071] Using the sucAB gene of *Enterobacter agglomerans* obtained as described above, a strain lacking α KGDH of *Enterobacter agglomerans* was obtained by homologous recombination.

[0072] First, pTWVEK101 was digested with *BglII* to remove the C-terminus region corresponding to about half of the sucA gene and the full length of the sucB gene. To this site, a chloramphenicol resistance gene fragment cut out from the pHSG399 (Takara Shuzo) with *AccI* was then inserted. The region of *sdhB- Δ sucAB::Cmr^r-sucC* obtained above was cut out with *AflIII* and *SacI*. The resulting DNA fragment was used to transform the *Enterobacter agglomerans*

AJ13355 strain by electroporation to obtain a chloramphenicol resistant strain, and thus a *Enterobacter agglomerans* AJ13356 strain lacking the *sucAB* gene where the *sucAB* gene on the chromosome was replaced by *sucAB::Cm^r* was obtained.

[0073] To confirm that the AJ13356 strain obtained as described above was deficient in the α KGDH activity, its enzymatic activity was determined by the method of Reed (L.J. Reed and B.B. Mukherjee, Methods in Enzymology 1969, 13, p.55-61). As a result, the α KGDH activity could not be detected in the AJ13356 strain as shown in Table 2, and thus it was confirmed that the strain lacked the *sucAB* as desired.

Table 2:

α KGDH activity	
Bacterial strain	α KGDH activity (Δ ABS/min/mg protein)
AJ13355	0.481
AJ13356	<0.0001

(5) Evaluation of L-glutamic acid productivity of *Enterobacter agglomerans* strain deficient in α KGDH

[0074] Each of the AJ13355 and AJ13356 strains was inoculated into a 500 ml-volume flask containing 20 ml of a culture medium comprising 40 g/L glucose, 20 g/L ammonium sulfate, 0.5 g/L magnesium sulfate heptahydrate, 2 g/L potassium dihydrogenphosphate, 0.5 g/L sodium chloride, 0.25 g/L calcium chloride heptahydrate, 0.02 g/L ferrous sulfate heptahydrate, 0.02 g/L manganese sulfate tetrahydrate, 0.72 mg/L zinc sulfate dihydrate, 0.64 mg/L copper sulfate pentahydrate, 0.72 mg/L cobalt chloride hexahydrate, 0.4 mg/L boric acid, 1.2 mg/L sodium molybdate dihydrate, 2 g/L yeast extract, 30 g/L calcium carbonate, 200 mg/L L-lysine monohydrochloride, 200 mg/L L-methionine and 200 mg/L DL- α,ϵ -diaminopimelic acid (DAP), and cultured at 37°C with shaking until the glucose contained in the culture medium was consumed. After the cultivation was completed, L-glutamic acid and α -ketoglutaric acid (abbreviated as " α KG" hereinafter) accumulated in the culture medium were measured. The results are shown in Table 3.

Table 3:

Accumulated amounts of L-glutamic acid and α KG		
Bacterial strain	Accumulated amount of L-glutamic acid	Accumulated amount of α KG
AJ13355	0.0 g/L	0.0 g/L
AJ13356	1.5	3.2

[0075] The AJ13356 strain deficient in the α KGDH activity accumulated 1.5 g/L of L-glutamic acid, and simultaneously accumulated 3.2 g/L of α KG.

(6) Introduction of RSFCPG into *Enterobacter agglomerans* strain lacking α KGDH and evaluation of L-glutamic acid productivity

[0076] The AJ13356 strain was transformed with the RSFCPG, and the resulting strain introduced with the RSFCPG, AJ13356/RSFCPG, was inoculated into a 500 ml-volume flask containing 20 ml of a culture medium comprising 40 g/L glucose, 20 g/L ammonium sulfate, 0.5 g/L magnesium sulfate heptahydrate, 2 g/L potassium dihydrogenphosphate, 0.5 g/L sodium chloride, 0.25 g/L calcium chloride heptahydrate, 0.02 g/L ferrous sulfate heptahydrate, 0.02 g/L manganese sulfate tetrahydrate, 0.72 mg/L zinc sulfate dihydrate, 0.64 mg/L copper sulfate pentahydrate, 0.72 mg/L cobalt chloride hexahydrate, 0.4 mg/L boric acid, 1.2 mg/L sodium molybdate dihydrate, 2 g/L yeast extract, 25 mg/L tetracycline, 30 g/L calcium carbonate, 200 mg/L L-lysine monohydrochloride, 200 mg/L L-methionine and 200 mg/L DL- α,ϵ -DAP, and cultured at 37°C with shaking until the glucose contained in the culture medium was consumed. After the cultivation was completed, L-glutamic acid and α KG accumulated in the culture medium were measured. The results are shown in Table 4.

Table 4:

Accumulated amounts of L-glutamic acid and α KG		
Bacterial strain	Accumulated amount of L-glutamic acid	Accumulated amount of α KG
AJ13356	1.4 g/L	2.9 g/L

EP 0 952 221 A2

Table 4: (continued)

Accumulated amounts of L-glutamic acid and α KG		
Bacterial strain	Accumulated amount of L-glutamic acid	Accumulated amount of α KG
AJ13356/RSFCPG	5.1	0.0

[0077] In the strain of which CS, PEPC and GDH activities were amplified by the introduction of RSFCPG, the accumulated amount of α KG was reduced, and the accumulated amount of L-glutamic acid was further improved.

Annex to the description

[0078]

5

10

15

20

25

30

35

40

45

50

55

EP 0 952 221 A2

SEQUENCE LISTING

5 <110> Ajinomoto Co., Inc.
 <120> L-Glutamic acid-producing bacterium and method for producing L-glutamic acid
 <130> EPA-53126
 <150> JP 10-69068
 10 <151> 1998-03-18
 <150> JP 10-297129
 <151> 1998-10-19
 <160> 7
 15 <170> PatentIn Ver. 2.0

 <210> 1
 <211> 4556
 <212> DNA
 20 <213> Enterobacter agglomerans
 <220>
 <221> CDS
 <222> (2)..(121)
 25 <220>
 <221> CDS
 <222> (322)..(3129)
 <220>
 30 <221> CDS
 <222> (3145)..(4368)
 <220>
 <221> CDS
 35 <222> (4437)..(4556)
 <400> 1
 t gca ttc agc gtt ttc cgc tgt cac agc atc atg aac tgt gta agt gtt 49
 Ala Phe Ser Val Phe Arg Cys His Ser Ile Met Asn Cys Val Ser Val
 1 5 10 15
 40 tgt cct aaa ggg cta aac ccg acg cgc gct atc ggc cac att aag tcg 97
 Cys Pro Lys Gly Leu Asn Pro Thr Arg Ala Ile Gly His Ile Lys Ser
 20 25 30
 atg ctg ctg caa cgc agc gcg tagttataacc accgggaacc tcaggttccc 148
 45 Met Leu Leu Gln Arg Ser Ala
 35
 ggtatitttac ggaagcctct gtaaagcgcg tcccaaccac gtttacaag gticccttac 208
 gggccgggacg cgcgctgcgc acagtgtcgc tatcgtgaa ctactacgc caaaccgcga 268
 50 aagcggcaac aaatgaaacc tcaaaaaagc ataacattgc ttaagggatc aca atg 324
 Met
 1
 cag aac agc gcg atg aag ccc tgg ctg gac tcc tcc tgg ctg gcc ggc 372
 55 Gln Asn Ser Ala Met Lys Pro Trp Leu Asp Ser Ser Trp Leu Ala Gly
 5 10 15

EP 0 952 221 A2

	gcg aat cag tct tac ata gag caa ctc tat gag gat ttc ctg acc gat	420
	Ala Asn Gln Ser Tyr Ile Glu Gln Leu Tyr Glu Asp Phe Leu Thr Asp	
	20 25 30	
5	cct gac tct gtg gat gca gtg tgg cgc tcg atg ttc caa cag tta cca	468
	Pro Asp Ser Val Asp Ala Val Trp Arg Ser Met Phe Gln Gln Leu Pro	
	35 40 45	
10	ggc acg gga gtg aaa cct gag cag ttc cac tcc gca act cgc gaa tat	516
	Gly Thr Gly Val Lys Pro Glu Gln Phe His Ser Ala Thr Arg Glu Tyr	
	50 55 60 65	
	ttc cgt cgc ctg gcg aaa gac gca tct cgt tac acc tcc tca gtt acc	564
	Phe Arg Arg Leu Ala Lys Asp Ala Ser Arg Tyr Thr Ser Ser Val Thr	
	70 75 80	
15	gat ccg gca acc aac tcc aaa caa gtg aaa gtg ctg cag ctg att aac	612
	Asp Pro Ala Thr Asn Ser Lys Gln Val Lys Val Leu Gln Leu Ile Asn	
	85 90 95	
20	gcg ttt cgt ttc cgc gga cat cag gaa gca aat ctc gat ccg ctt ggc	660
	Ala Phe Arg Phe Arg Gly His Gln Glu Ala Asn Leu Asp Pro Leu Gly	
	100 105 110	
	ctg tgg aaa cag gac cgc gtt gcc gat ctc gat cct gcc ttt cac gat	708
	Leu Trp Lys Gln Asp Arg Val Ala Asp Leu Asp Pro Ala Phe His Asp	
	115 120 125	
25	ctg acc gac gcc gat ttt cag gaa agc ttt aac gta ggt tct ttt gcc	756
	Leu Thr Asp Ala Asp Phe Gln Glu Ser Phe Asn Val Gly Ser Phe Ala	
	130 135 140 145	
30	att ggc aaa gaa acc atg aag ctg gcc gat ctg ttc gac gcg ctg aag	804
	Ile Gly Lys Glu Thr Met Lys Leu Ala Asp Leu Phe Asp Ala Leu Lys	
	150 155 160	
	cag acc tac tgt ggc tcg att ggt gca gag tat atg cac atc aat aac	852
	Gln Thr Tyr Cys Gly Ser Ile Gly Ala Glu Tyr Met His Ile Asn Asn	
	165 170 175	
	acc gaa gag aaa cgc tgg atc cag cag cgt atc gaa tcc ggt gcg agc	900
	Thr Glu Glu Lys Arg Trp Ile Gln Gln Arg Ile Glu Ser Gly Ala Ser	
	180 185 190	
40	cag acg tca ttc agt ggc gaa gag aaa aaa ggt ttc ctg aaa gag ctg	948
	Gln Thr Ser Phe Ser Gly Glu Glu Lys Lys Gly Phe Leu Lys Glu Leu	
	195 200 205	
	acc gcg gca gaa ggg ctg gaa aaa tat ctg ggc gcg aaa ttc ccg ggt	996
	Thr Ala Ala Glu Gly Leu Glu Lys Tyr Leu Gly Ala Lys Phe Pro Gly	
	210 215 220 225	
	gca aaa cgt ttc tcg ctg gaa ggc ggt gat gcg ctg gtg ccg atg ctg	1044
	Ala Lys Arg Phe Ser Leu Glu Gly Gly Asp Ala Leu Val Pro Met Leu	
	230 235 240	
50	gcg gag atg att cgt cat gcg ggc aaa agc ggc aca cgt gaa gtg gta	1092
	Arg Glu Met Ile Arg His Ala Gly Lys Ser Gly Thr Arg Glu Val Val	
	245 250 255	
55	ctg ggg atg gcg cac cgt ggc cgt ctt aac gta ctg att aac gta ctg	1140
	Leu Gly Met Ala His Arg Gly Arg Leu Asn Val Leu Ile Asn Val Leu	

EP 0 952 221 A2

	260	265	270	
	ggt aaa aag cca cag gat ctg ttc gac gaa ttc tcc ggt aaa cac aaa			1188
5	Gly Lys Lys Pro Gln Asp Leu Phe Asp Glu Phe Ser Gly Lys His Lys			
	275	280	285	
	gag cat ctg ggc acc ggt gat gtg aag tat cac atg ggc ttc tct tcg			1236
	Glu His Leu Gly Thr Gly Asp Val Lys Tyr His Met Gly Phe Ser Ser			
	290	295	300	305
10	gat att gaa acc gaa ggt ggt ctg gtg cat ctg gcg ctg gcg ttt aac			1284
	Asp Ile Glu Thr Glu Gly Gly Leu Val His Leu Ala Leu Ala Phe Asn			
	310	315	320	
15	ccg tct cac ctg gaa att gtc agc ccg gtg gtc atg gga tcg gta cgt			1332
	Pro Ser His Leu Glu Ile Val Ser Pro Val Val Met Gly Ser Val Arg			
	325	330	335	
	gca cgt ctc gat cgt ctg gcc gaa ccg gtc agc aat aaa gtg ttg cct			1380
	Ala Arg Leu Asp Arg Leu Ala Glu Pro Val Ser Asn Lys Val Leu Pro			
	340	345	350	
20	atc acc att cac ggt gat gcg gcg gtg att ggt cag ggc gtg gtt cag			1428
	Ile Thr Ile His Gly Asp Ala Ala Val Ile Gly Gln Gly Val Val Gln			
	355	360	365	
25	gaa acc ctg aac atg tct cag gcg cgc ggc tac gaa gtg ggc ggc acg			1476
	Glu Thr Leu Asn Met Ser Gln Ala Arg Gly Tyr Glu Val Gly Gly Thr			
	370	375	380	385
	gta cgt atc gtc att aac aac cag gtt ggt ttt acc acc tcc aac ccg			1524
	Val Arg Ile Val Ile Asn Asn Gln Val Gly Phe Thr Thr Ser Asn Pro			
	390	395	400	
30	aaa gat gcg cgt tca acc ccg tac tgt act gac atc ggc aag atg gtg			1572
	Lys Asp Ala Arg Ser Thr Pro Tyr Cys Thr Asp Ile Gly Lys Met Val			
	405	410	415	
35	ctg gca ccg att ttc cac gtc aat gct gac gat ccg gaa gcg gtg gcc			1620
	Leu Ala Pro Ile Phe His Val Asn Ala Asp Asp Pro Glu Ala Val Ala			
	420	425	430	
	ttt gtt acc cgc ctg gcg ctg gac tat cgc aac acc ttc aaa cgc gat			1668
	Phe Val Thr Arg Leu Ala Leu Asp Tyr Arg Asn Thr Phe Lys Arg Asp			
40	435	440	445	
	gtg ttt atc gat ctg gtg tgc tat cgc cgt cat ggt cac aac gag gcg			1716
	Val Phe Ile Asp Leu Val Cys Tyr Arg Arg His Gly His Asn Glu Ala			
	450	455	460	465
45	gat gag cca agt gct acc cag ccg ttg atg tac cag aaa atc aaa aag			1764
	Asp Glu Pro Ser Ala Thr Gln Pro Leu Met Tyr Gln Lys Ile Lys Lys			
	470	475	480	
50	cat ccg acg ccg cgt aaa att tac gcc gat cgt ctg gaa ggc gaa ggt			1812
	His Pro Thr Pro Arg Lys Ile Tyr Ala Asp Arg Leu Glu Gly Glu Gly			
	485	490	495	
	gtc gcg tcg cag gaa gat gcc acc gag atg gtg aac ctg tac cgc gat			1860
	Val Ala Ser Gln Glu Asp Ala Thr Glu Met Val Asn Leu Tyr Arg Asp			
	500	505	510	
55	gcg ctc gat gcg ggc gaa tgc gtg gtg ccg gaa tgg cgt ccg atg agc			1908

EP 0 952 221 A2

	Ala Leu Asp Ala Gly Glu Cys Val Val Pro Glu Trp Arg Pro Met Ser	
	515 520 525	
5	ctg cac tcc ttc acg tgg tgc cct tat ctg aac cac gaa tgg gat gag	1956
	Leu His Ser Phe Thr Trp Ser Pro Tyr Leu Asn His Glu Trp Asp Glu	
	530 535 540 545	
	cct tat ccg gca cag gtt gac atg aaa cgc ctg aag gaa ctg gca ttg	2004
	Pro Tyr Pro Ala Gln Val Asp Met Lys Arg Leu Lys Glu Leu Ala Leu	
10	550 555 560	
	cgt atc agc cag gtc cct gag cag att gaa gtg cag tgc cgc gtg gcc	2052
	Arg Ile Ser Gln Val Pro Glu Gln Ile Glu Val Gln Ser Arg Val Ala	
	565 570 575	
15	aag atc tat aac gat cgc aag ctg atg gcc gaa ggc gag aaa gcg ttc	2100
	Lys Ile Tyr Asn Asp Arg Lys Leu Met Ala Glu Gly Glu Lys Ala Phe	
	580 585 590	
	gac tgg ggc ggt gcc gag aat ctg gcg tac gcc acg ctg gtg gat gaa	2148
20	Asp Trp Gly Gly Ala Glu Asn Leu Ala Tyr Ala Thr Leu Val Asp Glu	
	595 600 605	
	ggt att ccg gtt cgc ctc tgc ggt gaa gac tcc ggt cgt gga acc ttc	2196
	Gly Ile Pro Val Arg Leu Ser Gly Glu Asp Ser Gly Arg Gly Thr Phe	
	610 615 620 625	
25	ttc cat cgc cac gcg gtc gtg cac aac cag gct aac ggt tca acc tat	2244
	Phe His Arg His Ala Val Val His Asn Gln Ala Asn Gly Ser Thr Tyr	
	630 635 640	
	acg ccg ctg cac cat att cat aac agc cag ggc gag ttc aaa gtc tgg	2292
30	Thr Pro Leu His His Ile His Asn Ser Gln Gly Glu Phe Lys Val Trp	
	645 650 655	
	gat tgc gtg ctg tct gaa gaa gcg gtg ctg gcg ttt gaa tac ggt tac	2340
	Asp Ser Val Leu Ser Glu Glu Ala Val Leu Ala Phe Glu Tyr Gly Tyr	
	660 665 670	
35	gcc acg gct gag ccg cgc gtg ctg acc atc tgg gaa gcg cag ttt ggt	2388
	Ala Thr Ala Glu Pro Arg Val Leu Thr Ile Trp Glu Ala Gln Phe Gly	
	675 680 685	
	gac ttt gcc aac ggt gct cag gtg gtg att gac cag ttc atc agc tct	2436
40	Asp Phe Ala Asn Gly Ala Gln Val Val Ile Asp Gln Phe Ile Ser Ser	
	690 695 700 705	
	ggc gaa cag aag tgg ggc cgt atg tgt ggc ctg gtg atg ttg ctg ccg	2484
	Gly Glu Gln Lys Trp Gly Arg Met Cys Gly Leu Val Met Leu Leu Pro	
45	710 715 720	
	cat ggc tac gaa ggt cag gga ccg gaa cac tcc tct gcc cgt ctg gaa	2532
	His Gly Tyr Glu Gly Gln Gly Pro Glu His Ser Ser Ala Arg Leu Glu	
	725 730 735	
50	cgc tat ctg caa ctt tgc gcc gag cag aac atg cag gtt tgc gtc ccg	2580
	Arg Tyr Leu Gln Leu Cys Ala Glu Gln Asn Met Gln Val Cys Val Pro	
	740 745 750	
	tcg acg ccg gct cag gtg tat cac atg ctg cgc cgt cag gcg ctg cgc	2628
55	Ser Thr Pro Ala Gln Val Tyr His Met Leu Arg Arg Gln Ala Leu Arg	
	755 760 765	

EP 0 952 221 A2

	ggg atg cgc cgt ccg ctg gtg gtg atg tcg ccg aag tcg ctg tta cgc	2676
	Gly Met Arg Arg Pro Leu Val Val Met Ser Pro Lys Ser Leu Leu Arg	
	770 775 780 785	
5	cat cca ctg gcg atc tcg tcg ctg gat gaa ctg gca aac ggc agt ttc	2724
	His Pro Leu Ala Ile Ser Ser Leu Asp Glu Leu Ala Asn Gly Ser Phe	
	790 795 800	
	cag ccg gcc att ggt gag atc gac gat ctg gat ccg cag ggc gtg aaa	2772
10	Gln Pro Ala Ile Gly Glu Ile Asp Asp Leu Asp Pro Gln Gly Val Lys	
	805 810 815	
	cgc gtc gtg ctg tgc tcc ggt aag gtt tac tac gat ctg ctg gaa cag	2820
	Arg Val Val Leu Cys Ser Gly Lys Val Tyr Tyr Asp Leu Leu Glu Gln	
	820 825 830	
15	cgt cgt aaa gac gag aaa acc gat gtt gcc atc gtg cgc atc gaa cag	2868
	Arg Arg Lys Asp Glu Lys Thr Asp Val Ala Ile Val Arg Ile Glu Gln	
	835 840 845	
	ctt tac ccg ttc ccg cat cag gcg gta cag gaa gca ttg aaa gcc tat	2916
20	Leu Tyr Pro Phe Pro His Gln Ala Val Gln Glu Ala Leu Lys Ala Tyr	
	850 855 860 865	
	tct cac gta cag gac ttt gtc tgg tgc cag gaa gag cct ctg aac cag	2964
	Ser His Val Gln Asp Phe Val Trp Cys Gln Glu Glu Pro Leu Asn Gln	
	870 875 880	
25	ggc gcc tgg tac tgt agc cag cat cat ttc cgt gat gtc gtg ccg ttt	3012
	Gly Ala Trp Tyr Cys Ser Gln His His Phe Arg Asp Val Val Pro Phe	
	885 890 895	
	ggg gcc acc ctg cgt tat gca ggt cgc ccg gca tcg gct tct ccg gcc	3060
30	Gly Ala Thr Leu Arg Tyr Ala Gly Arg Pro Ala Ser Ala Ser Pro Ala	
	900 905 910	
	gtg ggt tat atg tcc gta cac caa caa cag cag caa gac ctg gtt aat	3108
35	Val Gly Tyr Met Ser Val His Gln Gln Gln Gln Gln Asp Leu Val Asn	
	915 920 925	
	gac gca ctg aac gtc aat taattaaaag gaaagata atg agt agc gta gat	3159
	Asp Ala Leu Asn Val Asn Met Ser Ser Val Asp	
	930 935 1 5	
40	att ctc gtt ccc gac ctg cct gaa tcg gtt gca gat gcc aca gta gca	3207
	Ile Leu Val Pro Asp Leu Pro Glu Ser Val Ala Asp Ala Thr Val Ala	
	10 15 20	
	acc tgg cac aag aaa cca ggc gat gca gtc agc cgc gat gaa gtc atc	3255
45	Thr Trp His Lys Lys Pro Gly Asp Ala Val Ser Arg Asp Glu Val Ile	
	25 30 35	
	gtc gaa att gaa act gac aaa gtc gtg ctg gaa gtg ccg gca tct gcc	3303
	Val Glu Ile Glu Thr Asp Lys Val Val Leu Glu Val Pro Ala Ser Ala	
	40 45 50	
50	gat ggc gtg ctg gaa gcc gtg ctg gaa gac gaa ggg gca acc gtt acg	3351
	Asp Gly Val Leu Glu Ala Val Leu Glu Asp Glu Gly Ala Thr Val Thr	
	55 60 65	
55	tcc cgc cag atc ctg ggt cgc ctg aaa gaa ggc aac agt gcg ggt aaa	3399
	Ser Arg Gln Ile Leu Gly Arg Leu Lys Glu Gly Asn Ser Ala Gly Lys	

EP 0 952 221 A2

	70		75		80		85	
	gaa agc agt gcc aaa gcg gaa agc aat gac acc acg cca gcc cag cgt	3447						
5	Glu Ser Ser Ala Lys Ala Glu Ser Asn Asp Thr Thr Pro Ala Gln Arg							
	90		95		100			
	cag aca gcg tcg ctt gaa gaa gag agc agc gat gcg ctc agc ccg gcg	3495						
	Gln Thr Ala Ser Leu Glu Glu Glu Ser Ser Asp Ala Leu Ser Pro Ala							
	105		110		115			
10	atc cgt cgc ctg att gcg gag cat aat ctt gac gct gcg cag atc aaa	3543						
	Ile Arg Arg Leu Ile Ala Glu His Asn Leu Asp Ala Ala Gln Ile Lys							
	120		125		130			
	ggc acc ggc gta ggc gga cgt tta acg cgt gaa gac gtt gaa aaa cat	3591						
15	Gly Thr Gly Val Gly Gly Arg Leu Thr Arg Glu Asp Val Glu Lys His							
	135		140		145			
	ctg gcg aac aaa ccg cag gct gag aaa gcc gcc gcg cca gcg gcg ggt	3639						
	Leu Ala Asn Lys Pro Gln Ala Glu Lys Ala Ala Ala Pro Ala Ala Gly							
20	150		155		160		165	
	gca gca acg gct cag cag cct gtt gcc aac cgc agc gaa aaa cgt gtt	3687						
	Ala Ala Thr Ala Gln Gln Pro Val Ala Asn Arg Ser Glu Lys Arg Val							
	170		175		180			
	ccg atg acg cgt tta cgt aag cgc gtc gcg gag cgt ctg ctg gaa gcc	3735						
25	Pro Met Thr Arg Leu Arg Lys Arg Val Ala Glu Arg Leu Leu Glu Ala							
	185		190		195			
	aag aac agc acc gcc atg ttg acg acc ttc aac gaa atc aac atg aag	3783						
	Lys Asn Ser Thr Ala Met Leu Thr Thr Phe Asn Glu Ile Asn Met Lys							
30	200		205		210			
	ccg att atg gat ctg cgt aag cag tac ggc gat gcg ttc gag aag cgt	3831						
	Pro Ile Met Asp Leu Arg Lys Gln Tyr Gly Asp Ala Phe Glu Lys Arg							
	215		220		225			
35	cac ggt gtg cgt ctg ggc ttt atg tct ttc tac atc aag gcc gtg gtc	3879						
	His Gly Val Arg Leu Gly Phe Met Ser Phe Tyr Ile Lys Ala Val Val							
	230		235		240		245	
	gaa gcg ctg aag cgt tat cca gaa gtc aac gcc tct atc gat ggc gaa	3927						
40	Glu Ala Leu Lys Arg Tyr Pro Glu Val Asn Ala Ser Ile Asp Gly Glu							
	250		255		260			
	gac gtg gtg tac cac aac tat ttc gat gtg agt att gcc gtc tct acg	3975						
	Asp Val Val Tyr His Asn Tyr Phe Asp Val Ser Ile Ala Val Ser Thr							
	265		270		275			
45	cca cgc gga ctg gtg acg cct gtc ctg cgt gac gtt gat gcg ctg agc	4023						
	Pro Arg Gly Leu Val Thr Pro Val Leu Arg Asp Val Asp Ala Leu Ser							
	280		285		290			
	atg gct gac atc gag aag aaa att aaa gaa ctg gca gtg aaa ggc cgt	4071						
50	Met Ala Asp Ile Glu Lys Lys Ile Lys Glu Leu Ala Val Lys Gly Arg							
	295		300		305			
	gac ggc aag ctg acg gtt gac gat ctg acg ggc ggt aac ttt acc atc	4119						
	Asp Gly Lys Leu Thr Val Asp Asp Leu Thr Gly Gly Asn Phe Thr Ile							
	310		315		320		325	
55	acc aac ggt ggt gtg ttc ggt tcg ctg atg tct acg cca atc atc aac	4167						

EP 0 952 221 A2

Thr Asn Gly Gly Val Phe Gly Ser Leu Met Ser Thr Pro Ile Ile Asn
 330 335 340
 ccg cca cag agc gcg att ctg ggc atg cac gcc att aaa gat cgt cct 4215
 5 Pro Pro Gln Ser Ala Ile Leu Gly Met His Ala Ile Lys Asp Arg Pro
 345 350 355
 atg gcg gtc aat ggt cag gtt gtg atc ctg cca atg atg tac ctg gct 4263
 Met Ala Val Asn Gly Gln Val Val Ile Leu Pro Met Met Tyr Leu Ala
 360 365 370
 10 ctc tcc tac gat cac cgt tta atc gat ggt cgt gaa tct gtc ggc tat 4311
 Leu Ser Tyr Asp His Arg Leu Ile Asp Gly Arg Glu Ser Val Gly Tyr
 375 380 385
 15 ctg gtc gcg gtg aaa gag atg ctg gaa gat ccg gcg cgt ctg ctg ctg 4359
 Leu Val Ala Val Lys Glu Met Leu Glu Asp Pro Ala Arg Leu Leu Leu
 390 395 400 405
 gat gtc tgattcatca ctgggcacgc gtgctgtgcc caatctcaat actcttttca 4415
 Asp Val
 20
 gatctgaatg gatagaacat c atg aac tta cac gaa tac cag gct aaa cag 4466
 Met Asn Leu His Glu Tyr Gln Ala Lys Gln
 1 5 10
 25 ctg ttt gca cgg tat ggc atg cca gca ccg acc ggc tac gcc tgt act 4514
 Leu Phe Ala Arg Tyr Gly Met Pro Ala Pro Thr Gly Tyr Ala Cys Thr
 15 20 25
 aca cca cgt gaa gca gaa gaa gcg gca tcg aaa atc ggt gca 4556
 30 Thr Pro Arg Glu Ala Glu Glu Ala Ala Ser Lys Ile Gly Ala
 30 35 40
 <210> 2
 <211> 39
 35 <212> PRT
 <213> Enterobacter agglomerans
 <400> 2
 Ala Phe Ser Val Phe Arg Cys His Ser Ile Met Asn Cys Val Ser Val
 40 1 5 10 15
 Cys Pro Lys Gly Leu Asn Pro Thr Arg Ala Ile Gly His Ile Lys Ser
 20 25 30
 Met Leu Leu Gln Arg Ser Ala
 45 35
 <210> 3
 <211> 935
 50 <212> PRT
 <213> Enterobacter agglomerans
 <400> 3
 Met Gln Asn Ser Ala Met Lys Pro Trp Leu Asp Ser Ser Trp Leu Ala
 1 5 10 15
 55 Gly Ala Asn Gln Ser Tyr Ile Glu Gln Leu Tyr Glu Asp Phe Leu Thr

20					25					30						
Asp	Pro	Asp	Ser	Val	Asp	Ala	Val	Trp	Arg	Ser	Met	Phe	Gln	Gln	Leu	
35					40					45						
Pro	Gly	Thr	Gly	Val	Lys	Pro	Glu	Gln	Phe	His	Ser	Ala	Thr	Arg	Glu	
50					55					60						
Tyr	Phe	Arg	Arg	Leu	Ala	Lys	Asp	Ala	Ser	Arg	Tyr	Thr	Ser	Ser	Val	
65					70					75					80	
Thr	Asp	Pro	Ala	Thr	Asn	Ser	Lys	Gln	Val	Lys	Val	Leu	Gln	Leu	Ile	
85					90					95						
Asn	Ala	Phe	Arg	Phe	Arg	Gly	His	Gln	Glu	Ala	Asn	Leu	Asp	Pro	Leu	
100					105					110						
Gly	Leu	Trp	Lys	Gln	Asp	Arg	Val	Ala	Asp	Leu	Asp	Pro	Ala	Phe	His	
115					120					125						
Asp	Leu	Thr	Asp	Ala	Asp	Phe	Gln	Glu	Ser	Phe	Asn	Val	Gly	Ser	Phe	
130					135					140						
Ala	Ile	Gly	Lys	Glu	Thr	Met	Lys	Leu	Ala	Asp	Leu	Phe	Asp	Ala	Leu	
145					150					155					160	
Lys	Gln	Thr	Tyr	Cys	Gly	Ser	Ile	Gly	Ala	Glu	Tyr	Met	His	Ile	Asn	
165					170					175						
Asn	Thr	Glu	Glu	Lys	Arg	Trp	Ile	Gln	Gln	Arg	Ile	Glu	Ser	Gly	Ala	
180					185					190						
Ser	Gln	Thr	Ser	Phe	Ser	Gly	Glu	Glu	Lys	Lys	Gly	Phe	Leu	Lys	Glu	
195					200					205						
Leu	Thr	Ala	Ala	Glu	Gly	Leu	Glu	Lys	Tyr	Leu	Gly	Ala	Lys	Phe	Pro	
210					215					220						
Gly	Ala	Lys	Arg	Phe	Ser	Leu	Glu	Gly	Gly	Asp	Ala	Leu	Val	Pro	Met	
225					230					235					240	
Leu	Arg	Glu	Met	Ile	Arg	His	Ala	Gly	Lys	Ser	Gly	Thr	Arg	Glu	Val	
245					250					255						
Val	Leu	Gly	Met	Ala	His	Arg	Gly	Arg	Leu	Asn	Val	Leu	Ile	Asn	Val	
260					265					270						
Leu	Gly	Lys	Lys	Pro	Gln	Asp	Leu	Phe	Asp	Glu	Phe	Ser	Gly	Lys	His	
275					280					285						
Lys	Glu	His	Leu	Gly	Thr	Gly	Asp	Val	Lys	Tyr	His	Met	Gly	Phe	Ser	
290					295					300						
Ser	Asp	Ile	Glu	Thr	Glu	Gly	Gly	Leu	Val	His	Leu	Ala	Leu	Ala	Phe	
305					310					315					320	
Asn	Pro	Ser	His	Leu	Glu	Ile	Val	Ser	Pro	Val	Val	Met	Gly	Ser	Val	
325					330					335						
Arg	Ala	Arg	Leu	Asp	Arg	Leu	Ala	Glu	Pro	Val	Ser	Asn	Lys	Val	Leu	
340					345					350						
Pro	Ile	Thr	Ile	His	Gly	Asp	Ala	Ala	Val	Ile	Gly	Gln	Gly	Val	Val	
355					360					365						
Gln	Glu	Thr	Leu	Asn	Met	Ser	Gln	Ala	Arg	Gly	Tyr	Glu	Val	Gly	Gly	
370					375					380						
Thr	Val	Arg	Ile	Val	Ile	Asn	Asn	Gln	Val	Gly	Phe	Thr	Thr	Ser	Asn	
385					390					395					400	

EP 0 952 221 A2

	Pro	Lys	Asp	Ala	Arg	Ser	Thr	Pro	Tyr	Cys	Thr	Asp	Ile	Gly	Lys	Met
				405					410					415		
5	Val	Leu	Ala	Pro	Ile	Phe	His	Val	Asn	Ala	Asp	Asp	Pro	Glu	Ala	Val
			420						425				430			
	Ala	Phe	Val	Thr	Arg	Leu	Ala	Leu	Asp	Tyr	Arg	Asn	Thr	Phe	Lys	Arg
			435					440					445			
	Asp	Val	Phe	Ile	Asp	Leu	Val	Cys	Tyr	Arg	Arg	His	Gly	His	Asn	Glu
10		450					455					460				
	Ala	Asp	Glu	Pro	Ser	Ala	Thr	Gln	Pro	Leu	Met	Tyr	Gln	Lys	Ile	Lys
	465					470				475					480	
	Lys	His	Pro	Thr	Pro	Arg	Lys	Ile	Tyr	Ala	Asp	Arg	Leu	Glu	Gly	Glu
					485				490					495		
15	Gly	Val	Ala	Ser	Gln	Glu	Asp	Ala	Thr	Glu	Met	Val	Asn	Leu	Tyr	Arg
			500						505				510			
	Asp	Ala	Leu	Asp	Ala	Gly	Glu	Cys	Val	Val	Pro	Glu	Trp	Arg	Pro	Met
		515					520					525				
20	Ser	Leu	His	Ser	Phe	Thr	Trp	Ser	Pro	Tyr	Leu	Asn	His	Glu	Trp	Asp
		530					535				540					
	Glu	Pro	Tyr	Pro	Ala	Gln	Val	Asp	Met	Lys	Arg	Leu	Lys	Glu	Leu	Ala
	545					550				555					560	
25	Leu	Arg	Ile	Ser	Gln	Val	Pro	Glu	Gln	Ile	Glu	Val	Gln	Ser	Arg	Val
					565					570				575		
	Ala	Lys	Ile	Tyr	Asn	Asp	Arg	Lys	Leu	Met	Ala	Glu	Gly	Glu	Lys	Ala
			580						585				590			
30	Phe	Asp	Trp	Gly	Gly	Ala	Glu	Asn	Leu	Ala	Tyr	Ala	Thr	Leu	Val	Asp
		595						600				605				
	Glu	Gly	Ile	Pro	Val	Arg	Leu	Ser	Gly	Glu	Asp	Ser	Gly	Arg	Gly	Thr
		610					615				620					
35	Phe	Phe	His	Arg	His	Ala	Val	Val	His	Asn	Gln	Ala	Asn	Gly	Ser	Thr
	625					630					635				640	
	Tyr	Thr	Pro	Leu	His	His	Ile	His	Asn	Ser	Gln	Gly	Glu	Phe	Lys	Val
					645					650				655		
	Trp	Asp	Ser	Val	Leu	Ser	Glu	Glu	Ala	Val	Leu	Ala	Phe	Glu	Tyr	Gly
40			660						665				670			
	Tyr	Ala	Thr	Ala	Glu	Pro	Arg	Val	Leu	Thr	Ile	Trp	Glu	Ala	Gln	Phe
		675					680				685					
	Gly	Asp	Phe	Ala	Asn	Gly	Ala	Gln	Val	Val	Ile	Asp	Gln	Phe	Ile	Ser
45		690					695				700					
	Ser	Gly	Glu	Gln	Lys	Trp	Gly	Arg	Met	Cys	Gly	Leu	Val	Met	Leu	Leu
	705					710					715				720	
	Pro	His	Gly	Tyr	Glu	Gly	Gln	Gly	Pro	Glu	His	Ser	Ser	Ala	Arg	Leu
					725					730				735		
50	Glu	Arg	Tyr	Leu	Gln	Leu	Cys	Ala	Glu	Gln	Asn	Met	Gln	Val	Cys	Val
				740					745				750			
	Pro	Ser	Thr	Pro	Ala	Gln	Val	Tyr	His	Met	Leu	Arg	Arg	Gln	Ala	Leu
			755						760				765			
55	Arg	Gly	Met	Arg	Arg	Pro	Leu	Val	Val	Met	Ser	Pro	Lys	Ser	Leu	Leu

EP 0 952 221 A2

770 775 780
 Arg His Pro Leu Ala Ile Ser Ser Leu Asp Glu Leu Ala Asn Gly Ser
 785 790 795 800
 5 Phe Gln Pro Ala Ile Gly Glu Ile Asp Asp Leu Asp Pro Gln Gly Val
 805 810 815
 Lys Arg Val Val Leu Cys Ser Gly Lys Val Tyr Tyr Asp Leu Leu Glu
 820 825 830
 10 Gln Arg Arg Lys Asp Glu Lys Thr Asp Val Ala Ile Val Arg Ile Glu
 835 840 845
 Gln Leu Tyr Pro Phe Pro His Gln Ala Val Gln Glu Ala Leu Lys Ala
 850 855 860
 15 Tyr Ser His Val Gln Asp Phe Val Trp Cys Gln Glu Glu Pro Leu Asn
 865 870 875 880
 Gln Gly Ala Trp Tyr Cys Ser Gln His His Phe Arg Asp Val Val Pro
 885 890 895
 20 Phe Gly Ala Thr Leu Arg Tyr Ala Gly Arg Pro Ala Ser Ala Ser Pro
 900 905 910
 Ala Val Gly Tyr Met Ser Val His Gln Gln Gln Gln Gln Asp Leu Val
 915 920 925
 25 Asn Asp Ala Leu Asn Val Asn
 930 935

 <210> 4
 <211> 407
 30 <212> PRT
 <213> Enterobacter agglomerans
 <400> 4
 Met Ser Ser Val Asp Ile Leu Val Pro Asp Leu Pro Glu Ser Val Ala
 1 5 10 15
 35 Asp Ala Thr Val Ala Thr Trp His Lys Lys Pro Gly Asp Ala Val Ser
 20 25 30
 Arg Asp Glu Val Ile Val Glu Ile Glu Thr Asp Lys Val Val Leu Glu
 35 40 45
 40 Val Pro Ala Ser Ala Asp Gly Val Leu Glu Ala Val Leu Glu Asp Glu
 50 55 60
 Gly Ala Thr Val Thr Ser Arg Gln Ile Leu Gly Arg Leu Lys Glu Gly
 65 70 75 80
 45 Asn Ser Ala Gly Lys Glu Ser Ser Ala Lys Ala Glu Ser Asn Asp Thr
 85 90 95
 Thr Pro Ala Gln Arg Gln Thr Ala Ser Leu Glu Glu Glu Ser Ser Asp
 100 105 110
 50 Ala Leu Ser Pro Ala Ile Arg Arg Leu Ile Ala Glu His Asn Leu Asp
 115 120 125
 Ala Ala Gln Ile Lys Gly Thr Gly Val Gly Gly Arg Leu Thr Arg Glu
 130 135 140
 55 Asp Val Glu Lys His Leu Ala Asn Lys Pro Gln Ala Glu Lys Ala Ala
 145 150 155 160

EP 0 952 221 A2

Ala Pro Ala Ala Gly Ala Ala Thr Ala Gln Gln Pro Val Ala Asn Arg
165 170 175
5 Ser Glu Lys Arg Val Pro Met Thr Arg Leu Arg Lys Arg Val Ala Glu
180 185 190
Arg Leu Leu Glu Ala Lys Asn Ser Thr Ala Met Leu Thr Thr Phe Asn
195 200 205
10 Glu Ile Asn Met Lys Pro Ile Met Asp Leu Arg Lys Gln Tyr Gly Asp
210 215 220
Ala Phe Glu Lys Arg His Gly Val Arg Leu Gly Phe Met Ser Phe Tyr
225 230 235 240
15 Ile Lys Ala Val Val Glu Ala Leu Lys Arg Tyr Pro Glu Val Asn Ala
245 250 255
Ser Ile Asp Gly Glu Asp Val Val Tyr His Asn Tyr Phe Asp Val Ser
260 265 270
Ile Ala Val Ser Thr Pro Arg Gly Leu Val Thr Pro Val Leu Arg Asp
275 280 285
20 Val Asp Ala Leu Ser Met Ala Asp Ile Glu Lys Lys Ile Lys Glu Leu
290 295 300
Ala Val Lys Gly Arg Asp Gly Lys Leu Thr Val Asp Asp Leu Thr Gly
305 310 315 320
25 Gly Asn Phe Thr Ile Thr Asn Gly Gly Val Phe Gly Ser Leu Met Ser
325 330 335
Thr Pro Ile Ile Asn Pro Pro Gln Ser Ala Ile Leu Gly Met His Ala
340 345 350
30 Ile Lys Asp Arg Pro Met Ala Val Asn Gly Gln Val Val Ile Leu Pro
355 360 365
Met Met Tyr Leu Ala Leu Ser Tyr Asp His Arg Leu Ile Asp Gly Arg
370 375 380
35 Glu Ser Val Gly Tyr Leu Val Ala Val Lys Glu Met Leu Glu Asp Pro
385 390 395 400
Ala Arg Leu Leu Leu Asp Val
405

40 <210> 5
<211> 40
<212> PRT
<213> Enterobacter agglomerans
45 <400> 5
Met Asn Leu His Glu Tyr Gln Ala Lys Gln Leu Phe Ala Arg Tyr Gly
1 5 10 15
Met Pro Ala Pro Thr Gly Tyr Ala Cys Thr Thr Pro Arg Glu Ala Glu
20 25 30
50 Glu Ala Ala Ser Lys Ile Gly Ala
35 40
55 <210> 6
<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 6

gtcgacaata gccygaatct gttctggtcg

30

<210> 7

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 7

aagcttatcg acgctcccct ccccaccgtt

30

Claims

1. A microorganism belonging to the genus *Enterobacter* or *Serratia* having the ability to produce L-glutamic acid and having at least one of the following properties:

(a) the microorganism has increased activity of an enzyme catalyzing a reaction in the L-glutamic acid biosynthesis; and

(b) the microorganism has decreased activity or is deficient in an activity of an enzyme catalyzing a reaction branching from the pathway of the L-glutamic acid biosynthesis and producing a compound other than L-glutamic acid.

2. A microorganism according to claim 1 wherein the enzyme catalyzing the reaction in the L-glutamic acid biosynthesis is at least one selected from the group consisting of citrate synthase, phosphoenolpyruvate carboxylase, and glutamate dehydrogenase.

3. A microorganism according to claim 2 wherein the enzyme catalyzing the reaction in the L-glutamic acid biosynthesis includes all of citrate synthase, phosphoenolpyruvate carboxylase, and glutamate dehydrogenase.

4. A microorganism according to any one of claims 1 to 3 wherein the enzyme catalyzing the reaction branching from the pathway of the L-glutamic acid biosynthesis and producing a compound other than L-glutamic acid is α -ketoglutarate dehydrogenase.

5. A microorganism according to any one of claims 1 to 4 which is *Enterobacter agglomerans* or *Serratia liquefaciens*.

6. A method for producing L-glutamic acid which comprises culturing the microorganism as defined in any one of claims 1 to 5 in a liquid culture medium to produce and accumulate L-glutamic acid in the culture medium, and collecting the L-glutamic acid from the culture medium.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

[88.0% / 935 aa]

```

1'  MQNSAMKPYLDSSWLAGANQSYIEQLYEDFLTPDSDAVVRSMFQQLPGTGVKPEQFHS
1"  MQNSALKAWLDSSYLSGANQSWIEQLYEDFLTPDSDANWRSTFQQLPGTGVKPDQFHS
61'  ATREYFRRLAKDASRYTSSVTPATNSKQVKVLQLINAFRFRGHQEANLDPLGLWKQDRV
61"  QTREYFRRLAKDASRYSSITSDPDTHVKQVKVLQLINAYRFRGHQHANLDPLGLWQQQKV
121' ADLDPAFHDLTDADFQESFNVGSFAIGKETMKLADLFDALKQTYCGSIGAEYMHINTEE
121" ADLDPSEHDLTEADFQETFNVGSFASGKETMKLGELLEALKQTYCGPIGAEYMHITSTEE
181' KRWIQQRIESGASQTSFSGEEKKGFLKELTAAEGLEKYLGAKEFGAKRFSLEGGDALVPM
181" KRWIQQRIESG--RATFNSEKKRFLSELTAAEGLERYLGAKEFGAKRFSLEGGDALIPM
241' LREMIRHAGKSGTREVVLQMAHRGRLNVLINVLGKKPQDLDEFSGKHKEHLGTGOVKYH
239" LKEMIRHAGNSGTREVVLQMAHRGRLNVLNVLGKKPQDLDEFAGKHKEHLGTGOVKYH
301' MGFSSDIETEGGLVHLALAFNPShLEIVSPVVMGsvRARLDRLAEPVSNKVLpITIHGDA
299" MGFSSDFQTDGGLVHLALAFNPShLEIVSPVVGsvRARLDRLDEPSSNKVLpITIHGDA
361' AVIGQGVVQETLNMSQARGYEVGGTVRIVINNQGFTTsnPKDARSTpyCTDIGKmvLAP
359" AVTGQGVVQETLNMSKARGYEVGGTVRIVINNQGFTTsnPLDARSTpyCTDIGKmvQAP
421' IFHVNADDPEAVAFVTRLALDYRNTFKRDVFIDLVCYRRHGHNEADEPSATQPLMYQKIK
419" IFHVNADDPEAVAFVTRLALDFRNTFKRDVFIDLVSYRRHGHNEADEPSATQPLMYQKIK
481' KHPTPRKIYADRLEGEVSAQEDATEMVNLYRDALDAGECVVPEWRPMSLHSFTWSPYLN
479" KHPTPRKIYADKLEQEKVATLEDATEMVNLYRDALDAGDCVVAEWRPMNMHSFTWSPYLN
541' HEWDEPYPAQVDMKRLKELALRISQVPEQIEVQSRVAKIYNDRKLMAEGEKAFDWGGAEN
539" HEWDEEYPNKVEMKRLQELAKRISTVPEAVEMQSRVAKIYGDRQAMAAGEKLFDWGGAEN
601' LAYATLVDEGIPVRLSGEDSGRGTFFHRHAVVHNQANGSTYTPLHHIHNSQGEFKVWDSV
599" LAYATLVDEGIPVRLSGEDSGRGTFFHRHAVIHNSNGSTYTPLQHHHNGQGAFRVWDSV
661' LSEEAVLAFEYGYATAEPRVLTIWEAQFGDFANGAQVVIDQFISSGEQKWGRMCGLVMLL
659" LSEEAVLAFEYGYATAEPRTLTIWEAQFGDFANGAQVVIDQFISSGEQKWGRMCGLVMLL
721' PHGYEGQGPEHSSARLERYLQLCAEQNMQVCVPSTPAQVYHMLRRQALRGMRRLVVMSP
719" PHGYEGQGPEHSSARLERYLQLCAEQNMQVCVPSTPAQVYHMLRRQALRGMRRLVVMSP
781' KSLLRHPLAISSLELANGSFQPAIGEIDDLDPQGVKRVLCSGKVYYDLLEQRRKDEKT
779" KSLLRHPLAVSSLEELANGTFLPAIGEIDELDPKGVKRVMCSGKVYYDLLEQRRKNQH
841' DVAIVRIEQLYFPFHQAVQEALKAYSHVQDFVYCQEEPLNQGAWYCSQHHFRDVPFGAT
839" DVAIVRIEQLYFPFKAMQEVLLQFAHYKDFVYCQEEPLNQGAWYCSQHHFREVIPFGAS
901' LRYAGRPASASPAVGYSVHQQQQDLVNDALNVN
899" LRYAGRPASASPAVGYSVHQQQQDLVNDALNVE

```

FIG. 7

[88.2% / 407 aa]

```

1' MSSVDILVPDLPE$VADATVATWHKKPGDAVSRDEVIVEIETDKVVLEVPASADGVLEAV
.....
1" MSSVDILVPDLPE$VADATVATWHKKPGDAVVRDEVLVEIETDKVVLEVPASADGILDAV

61' LEDEGATVTSRQILGRLKEGNSAGKES$AKAESNDTTPAQRTASLEEESDAL$PAIRR
.....
61" LEDEGTTVTSRQILGRLREGNSAGKET$AKSEEKASTPAQRQQASLEEQNNDAL$PAIRR

121' LIAEHNLDAAQIKGTGVGGRLTREDVEKHLANKPQAEKAAAPAAGAATAQQPVANRSEKR
.....
121" LLAEHNLDASAIKGTGVGGRLTREDVEKHLAKAPAKE--SAPAAAAAPAAQPALAARSEKR

181' VPMTRLRKRAERLLEAKNSTAMLTTFNEINMKPIMDLRKQYGDAFEKRHGVRLGFMSFY
.....
179" VPMTRLRKRAERLLEAKNSTAMLTTFNEVNMKPIMDLRKQYGEAFEKRHGIRLGFMSFY

241' IKAVVEALKRYPEVNASIDGEDVYHNYFDVSI$AVSTPRGLVTPVLRD$DAL$SMADIEKK
.....
239" VKAVVEALKRYPEVNASIDGDDVYHNYFDVSM$AVSTPRGLVTPVLRD$DTL$GMADIEKK

301' IKELAVKGRDGKLTVDLTTGGNFTITNGGVFGSLMSTPIINPPQSAILGMHAIKDRPMAV
.....
299" IKELAVKGRDGKLTVEDLTGGNFTITNGGVFGSLMSTPIINPPQSAILGMHAIKDRPMAV

361' NGQVVILPMMYLALSYDHRLIDGRESVG$YLVAVKEMLEDPARLLLDV
.....
359" NGQVEILPMMYLALSYDHRLIDGRESVG$FLVTIKELLEDPTRLLLDV

```

FIG. 8

[95.1% / 41 aa]

```

1' MNLHEYQAKQLFARYGMPAPTGYACTTPREAE$EAASKIGAG
.....
1" MNLHEYQAKQLFARYGLPAPVGYACTTPREAE$EAASKIGAGPMVVKQVHAGGRGKAGGV

```

FIG. 9

[97.4% / 39 aa]

```

1' AF$VFRCHSIMNCV$VCPKGLNPTRAIGHIK$MLLQRS
.....
181" FLIDSRDTETDSRLDGLSDAF$VFRCHSIMNCV$VCPKGLNPTRAIGHIK$MLLQRN

```

FIG. 10

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 952 221 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:

12.09.2001 Bulletin 2001/37

(43) Date of publication A2:

27.10.1999 Bulletin 1999/43

(21) Application number: 99105508.8

(22) Date of filing: 17.03.1999

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 18.03.1998 JP 6906898

19.10.1998 JP 29712998

(71) Applicant: Ajinomoto Co., Inc.

Tokyo (JP)

(72) Inventors:

- Izui, Hiroshi c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)

- Ono, Eiji c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Matsui, Kazuhiko c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Moriya, Mika c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Ito, Hisao c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)
- Hara, Yoshihiko c/o Ajinomoto Co., Inc.
Kawasaki-ku, Kawasaki-shi, Kanagawa-ken (JP)

(74) Representative: Strehl Schübel-Hopf & Partner
Maximilianstrasse 54
80538 München (DE)

(54) **L-glutamic acid-producing bacterium and method for producing L-glutamic acid**

(57) L-Glutamic acid is produced by culturing in a liquid culture medium a microorganism belonging to the genus *Enterobacter* or *Serratia* and having an ability to produce L-glutamic acid, which increases in an activity of enzyme catalyzing a reaction for L-glutamic acid bio-

synthesis, or which decreases in or is deficient in an activity of an enzyme catalyzing a reaction branching from a pathway for L-glutamic acid biosynthesis and producing a compound other than L-glutamic acid, and collecting produced L-glutamic acid from the culture medium.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 10 5508

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X, D	EP 0 670 370 A (AJINOMOTO KK) 6 September 1995 (1995-09-06) * the whole document *	1, 2, 4, 6	C12N15/53 C12N15/60 C12N1/21 C12N1/20
X	WO 97 08294 A (AJINOMOTO KK ; MATSUI KAZUHIKO (JP); HUKASE KUMIKO (JP); TSUJIMOTO) 6 March 1997 (1997-03-06) * the whole document *	1, 2, 4, 6	C12P13/14 //(C12N1/20, C12R1:01, 1:425)
E	EP 1 078 989 A (AJINOMOTO KK) 28 February 2001 (2001-02-28) * the whole document *	1-6	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C12N C12P
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 20 July 2001	Examiner Oderwald, H
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 10 5508

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-07-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0670370 A	06-09-1995	JP 7203980 A	08-08-1995
		BR 9500052 A	03-10-1995
		CN 1128295 A	07-08-1996
		US 5573945 A	12-11-1996
WO 9708294 A	06-03-1997	BR 9610016 A	06-07-1999
		CN 1193343 A	16-09-1998
		DE 19681532 T	03-12-1998
		FR 2738014 A	28-02-1997
		US 6110714 A	29-08-2000
EP 1078989 A	28-02-2001	AU 5345200 A	22-02-2001
		BR 0003695 A	05-06-2001
		CN 1292421 A	25-04-2001
		PL 341974 A	26-02-2001